arXiv:hep-ex/9707003v1 1 Jul 1997

UCL/HEP 97-03

ENERGY FLOWS AND JET PRODUCTION IN TAGGED $e\gamma$ EVENTS AT LEP

A.M. ROOKE FOR THE OPAL COLLABORATION
Department of Physics, UCL, Gower Street, London WC1E 6BT, UK.

1 Introduction

It has been shown [1] that the predictions of the hadronic energy flow in $e\gamma$ DIS processes by the QCD-based Monte Carlo generators, HERWIG and PYTHIA, disagree with the data in the $Q^2 = 6$–$30\ \mathrm{GeV}^2$ region. The disagreements are particularly marked in the regions of pseudorapidity well measured by the OPAL detector and hence where jets can be accurately reconstructed.

	0 jet	1 jet	2 jet
Data	30.7%	63.8%	5.4%
HERWIG	34.0%	63.6%	2.4%
PYTHIA	32.8%	65.5%	1.7%

Table 1: Jet rates of HERWIG and PYTHIA compared to the data.

2 Results And Conclusions

Table 1 shows that the fraction of data events with 2 cone jets (of transverse energy $\mathrm{E}_{\mathrm{T,jet}} > 3\mathrm{GeV}$ and pseudorapidity $|\eta_{\mathrm{jet}}| < 2$) exceeds predictions from the HERWIG and PYTHIA samples by over a factor of 2. Fig. 1a) shows the average hadronic energy flow per event as a function of pseudorapidity and b), c) and d) show the contributions to a) from events with 0, 1 or 2 jets. The underestimation by HERWIG and PYTHIA of events with 2 jets combined with their imperfect modelling of energy flow (particularly in the pseudorapidity region $0.9 < \eta < 1.8$ for events with 1 jet) give rise to the previously reported discrepancies [1] shown in fig. 1a). By selecting events with 2 identified hadron cone jets, we have found a subset of the data in which disagreements with Monte Carlo predictions are greatly enhanced. This result has been used to constrain improvements to the models [2].



Figure 1: Average energy flow in an event as a function of pseudorapidity $\eta = -\ln\tan\frac{\theta}{2}$

References

1. OPAL Collab., K. Ackerstaff *et al*, *Z. Phys.* C **74**, 33 (1997)
2. J.A. Lauber, L. Lönnblad and M.H. Seymour, *these proceedings*